



Contribution ID : 99

Type : **Poster**

## Using synchrotron radiation to determine the X-ray structure and ct-DNA binding affinity of platinum(II) anticancer complexes

*Thursday, 20 November 2014 17:30 (90)*

Platinum(II) anticancer complexes incorporating 2,2'-bipyridine (bpy), 4,4'-dimethyl-2,2'-bipyridine (44Me2bpy) and 2-(2'-pyridyl)quinoxaline (2pq) as polyaromatic ligands and cyclic diamines as ancillary ligands have been synthesised and were characterised via several methods including synchrotron radiation X-ray crystallography. The crystal structure of  $[\text{Pt}(44\text{Me2bpy})(1\text{S},2\text{S-diaminocyclohexane})]^{2+}$  (44MEBSS) revealed a square planar coordination geometry similar to other complexes of this type whereas the complex  $[\text{Pt}(2\text{pq})(1\text{S},2\text{S-diaminocyclohexane})]^{2+}$  (2PQSS) was distorted square planar. The binding of 2PQSS and 44MEBSS to calf-thymus DNA (ct-DNA) was analysed using synchrotron radiation circular dichroism (SRCD) melting experiments and compared to similar complexes that incorporate 1,10-phenanthroline and dipyrdo[3,2-f:2',3'-h]quinoxaline. The results revealed unexpected trends in DNA affinity relative to polyaromatic ligand size.

### Keywords or phrases (comma separated)

synchrotron radiation circular dichroism (SRCD) melting experiments

### Summary

**Primary author(s) :** Mr PAGES, Benjamin (University of Western Sydney)

**Co-author(s) :** Dr LI, Feng (University of Western Sydney); Prof. ALDRICH-WRIGHT, Janice (University of Western Sydney); Dr JONES, Nikola (Department of Physics and Astronomy - Institute for Storage Ring Facilities, Aarhus University, Denmark); Dr ZHANG, Yingjie Zhang (ANSTO)

**Presenter(s) :** Mr PAGES, Benjamin (University of Western Sydney)

**Session Classification :** Welcome Function, Poster Session, Exhibition

**Track Classification :** Biological Systems