



# USER MEETING 2016

24-25 NOVEMBER

National Centre for Synchrotron Science



Ansto

Australian Synchrotron

Contribution ID : 226

Type : Poster

## Ni/TiO<sub>2</sub> - Low Cost Photocatalysts for Solar H<sub>2</sub> Production

This work targets the development of efficient metal co-catalyst modified titania photocatalysts for alcohol photoreforming to H<sub>2</sub> that function under direct sunlight. Conventionally, noble metals such as platinum, palladium or gold have been used co-catalysts to activate TiO<sub>2</sub> for hydrogen production, though the use of such co-catalysts for industrial scale H<sub>2</sub> manufacture is not feasible due to their high cost and low natural abundance, motivating the search for low cost alternatives.

This study compares the performance of 3 different Ni/TiO<sub>2</sub> photocatalysts for H<sub>2</sub> production in alcohol-water mixtures, placing particular emphasis on the role of the TiO<sub>2</sub> support and alcohol sacrificial reagent. P25 TiO<sub>2</sub> (85% Anatase, 15% Rutile), isolate anatase from P25 TiO<sub>2</sub>, isolate rutile from P25 TiO<sub>2</sub>, commercial brookite and physical mixed P25 TiO<sub>2</sub> were used as the support phase. XPS and Ni L-edge NEXAFS analyses verified that metallic Ni was the dominant nickel species in the near surface region of the photocatalysts. Ti L-edge NEXAFS spectra show L3-edge and L2-edge features with two sublevels (t<sub>2g</sub> and e<sub>g</sub>) which arise from crystal-field splitting caused by the octahedral ligand fields about the Ti<sup>4+</sup> cations. The difference between the three TiO<sub>2</sub> polymorphs (anatase, rutile and brookite) was seen in the e<sub>g</sub> feature at the L3 edge, which is split into two unresolved components (i.e. d<sub>z<sup>2</sup></sub> and d<sub>x<sup>2</sup>-y<sup>2</sup></sub> states). This change in the relative intensity of the d<sub>z<sup>2</sup></sub> and d<sub>x<sup>2</sup>-y<sup>2</sup></sub> peaks on going from anatase to rutile is explained by distortion of the Ti<sup>4+</sup> site from D<sub>2h</sub> (in anatase) to D<sub>2d</sub> (in rutile). For brookite, the d<sub>z<sup>2</sup></sub> and d<sub>x<sup>2</sup>-y<sup>2</sup></sub> peaks have similar intensities. These observations are in good accord with Ti L-edge data reported for mineral and synthetic titanias (Fig. 1). The Ni/P25 TiO<sub>2</sub> photocatalysts were very active for H<sub>2</sub> production in 10 vol.% alcohol-water mixtures under UV excitation, with the optimal Ni loading being ~0.5 wt.%. Ni/anatase and Ni/physical mixed P25 photocatalysts showed a diminution in the photocatalytic H<sub>2</sub> production performance, which confirmed the importance of interfacial electron transfer at the rutile:anatase interface.

Indico rendering error

Could not include image: Problem downloading image (<http://noigraphics.heliohost.org/fig1.png>)

### Keywords or phrases (comma separated)

H<sub>2</sub> production; Ni/TiO<sub>2</sub>

### Are you a student?

Yes

### Do you wish to take part in the Student Poster Slam?

No

### Are you an ECR? (<5 yrs since PhD/Masters)

No

## **What is your gender?**

Female

**Primary author(s):** Ms CHEN, Wan-Ting (The University of Auckland)

**Co-author(s):** Dr WATERHOUSE, Geoffrey I.N. (The University of Auckland)

**Presenter(s):** Ms CHEN, Wan-Ting (The University of Auckland)

**Track Classification :** Advanced Materials