



Contribution ID : 150

Type : Oral

With the synchrotron and beyond the synchrotron towards the structure of the TIR-domain signallosome

Friday, 25 November 2016 15:15 (50)

TIR (Toll/interleukin-1 receptor, resistance protein) domains are key components of innate immunity signaling pathways. They are found in animals, plants and bacteria, for example in TLRs (Toll-like receptors) and TLR adaptors in animals, NLRs (nucleotide binding, leucine-rich repeat receptors) in plants, and virulence factors interfering with immune responses in bacteria. While it has been well established that signaling depends on regulated self-association and homotypic association of TIR domains, every single TIR domain structure has revealed a different association mode [1]. In the search for common features, we have targeted a number of TIR domains from mammals, plants and bacteria to characterize structurally. We used the Australian Synchrotron to determine a number of TIR-domain crystal structures and study association using SAXS, including those from the human TLR adaptor proteins MAL [2] and SARM (unpublished), the bacterial protein TcpB from Brucella melitensis [3] and the plant immune proteins L6 from flax [4], RPS4 and RRS1 from Arabidopsis [5], SNC1 and RPP1 from Arabidopsis and MrRPV1 from grapevine (unpublished). These crystal structures have started revealing common trends in the TIR-domain association modes, in particular for bacterial and plant TIR domains. Furthermore, for the TLR adaptors MAL and MyD88, we have been able to reconstitute large assemblies and determine the structure for the former by cryo-electron microscopy (unpublished), while we are characterizing the structure of the latter by synchrotron and X-FEL-based serial crystallography. Jointly, these studies suggest a general mechanism of function of TIR domains, which involves "signalling by cooperative assembly formation (SCAF)" with prion-like features that is consistent with signaling in other innate immunity pathways.

References

[1] Ve et al, Structure and function of Toll/interleukin-1 receptor/resistance protein (TIR) domains, Apoptosis, 20 (2015), 250-61

[2] Valkov et al, "Crystal structure of TLR adaptor MAL/TIRAP reveals the molecular basis for signal transduction and disease protection", Proc Natl Acad Sci USA, 108 (2011), 14879-14884

[3] Alaidarous et al, Mechanism of bacterial interference with TLR4 signaling by Brucella TIR-domain-containing protein TcpB, J Biol Chem, 289 (2014), 654-68

[4] Bernoux et al, Structural and functional analysis of a plant resistance protein TIR domain reveals interfaces for self-association, signaling and autoregulation, Cell Host Microbe, 9 (2011), 200-211

Williams et al, Structural basis for assembly and function of a heterodimeric plant immune receptor, Science 344 (2014), 299-303

Keywords or phrases (comma separated)

crystallography, SAXS, higher-order assembly signalling

Are you a student?

No

Do you wish to take part in</br>the Student Poster Slam?

No

Are you an ECR? (<5 yrs</br>since PhD/Masters)

No

What is your gender?

Male

Primary author(s): Prof. KOBE, Bostjan (University of Queensland)Presenter(s): Prof. KOBE, Bostjan (University of Queensland)Session Classification: Plenary 3

Track Classification : Structural Biology